

The Container Store

EXTRANEAL Patient Kit

Important Safety Information about EXTRANEAL (Icodextrin)

Whenever you receive medical care - whether it's a scheduled appointment, an outpatient procedure or an emergency room visit - be sure to bring your **EXTRANEAL** patient kit along with you.

Tell the doctors and nurses that you are using **EXTRANEAL**, and that some glucose monitors and test strips may give a false high glucose reading.

This kit has additional information related to glucose monitors for nurses and doctors.

SC91118-002

December

Dear Doctor,

Extraneal (Icodextrin) is a solution used for peritoneal dialysis. It is important to be aware of the following information:

Extraneal (Icodextrin) may cause hypoglycemia. Patients may experience symptoms of low blood sugar, such as dizziness, weakness, and confusion. Patients should be aware of these symptoms and should be treated accordingly.

ONLY use the EXTRANEAL test strips and additional information provided in this kit.

The Extraneal (Icodextrin) solution may contain small amounts of glucose. This glucose should be taken into account when calculating total glucose intake.

DO NOT use Extraneal (Icodextrin) if you are allergic to icodextrin or any of the other ingredients. In addition, if you are taking other medications, you may experience hypoglycemia. Extraneal (Icodextrin) may mask the effects of hypoglycemia. Patients should be aware of these symptoms and should be treated accordingly.

Additional information is provided in the EXTRANEAL patient kit. Please read this information carefully. For more information, please contact your healthcare provider.

1. Check for the EXTRANEAL test strips.

Information is provided in the EXTRANEAL patient kit. Please read this information carefully.

Extraneal (Icodextrin) may cause hypoglycemia. Patients may experience symptoms of low blood sugar, such as dizziness, weakness, and confusion. Patients should be aware of these symptoms and should be treated accordingly.

ONLY use the EXTRANEAL test strips and additional information provided in this kit.

The Extraneal (Icodextrin) solution may contain small amounts of glucose. This glucose should be taken into account when calculating total glucose intake.

DO NOT use Extraneal (Icodextrin) if you are allergic to icodextrin or any of the other ingredients. In addition, if you are taking other medications, you may experience hypoglycemia. Extraneal (Icodextrin) may mask the effects of hypoglycemia. Patients should be aware of these symptoms and should be treated accordingly.

1. Check for the EXTRANEAL test strips.